


6. Large deviations


6. Large deviations



6. Large deviations

Characterized by

$$\Lambda_{N,t}(x) := -N^{-1} \log \mathbb{P}_{\psi_{N,t}} \left[N^{-1} \sum_{i=1}^N (Y_i^{N,t} - \langle \varphi_t, O\varphi_t \rangle) > x \right] .$$



6. Large deviations

Characterized by

$$\Lambda_{N,t}(x) := -N^{-1} \log \mathbb{P}_{\psi_{N,t}} \left[N^{-1} \sum_{i=1}^N (Y_i^{N,t} - \langle \varphi_t, O\varphi_t \rangle) > x \right] .$$

Kirkpatrick-R.-Schlein '20, R.-Seiringer '21: There exists, $C_{1,t}, C_{2,t} > 0$ s.t.

$$\limsup_{N \rightarrow \infty} \Lambda_{N,t}(x) \leq \frac{x^2}{2\sigma_t^2} + C_{1,t}x^3, \quad \liminf_{N \rightarrow \infty} \Lambda_{N,t}(x) \geq \frac{x^2}{2\sigma_t^2} - C_{2,t}x^{5/2}$$


6. Large deviations

Characterized by

$$\Lambda_{N,t}(x) := -N^{-1} \log \mathbb{P}_{\psi_{N,t}} \left[N^{-1} \sum_{i=1}^N (Y_i^{N,t} - \langle \varphi_t, O \varphi_t \rangle) > x \right].$$

Kirkpatrick-R.-Schlein '20, R.-Seiringer '21: There exists, $C_{1,t}, C_{2,t} > 0$ s.t.

$$\limsup_{N \rightarrow \infty} \Lambda_{N,t}(x) \leq \frac{x^2}{2\sigma_t^2} + C_{1,t}x^3, \quad \liminf_{N \rightarrow \infty} \Lambda_{N,t}(x) \geq \frac{x^2}{2\sigma_t^2} - C_{2,t}x^{5/2}$$



6. Large deviations


Characterized by

$$\Lambda_{N,t}(x) := -N^{-1} \log \mathbb{P}_{\psi_{N,t}} \left[N^{-1} \sum_{i=1}^N (Y_i^{N,t} - \langle \varphi_t, O\varphi_t \rangle) > x \right].$$

Kirkpatrick-R.-Schlein '20, R.-Seiringer '21: There exists, $C_{1,t}, C_{2,t} > 0$ s.t.

$$\limsup_{N \rightarrow \infty} \Lambda_{N,t}(x) \leq \frac{x^2}{2\sigma_t^2} + C_{1,t}x^3, \quad \liminf_{N \rightarrow \infty} \Lambda_{N,t}(x) \geq \frac{x^2}{2\sigma_t^2} - C_{2,t}x^{5/2}$$

► Variance σ_t^2 agrees with CLTs



6. Large deviations

Characterized by


$$\Lambda_{N,t}(x) := -N^{-1} \log \mathbb{P}_{\psi_{N,t}} \left[N^{-1} \sum_{i=1}^N (Y_i^{N,t} - \langle \varphi_t, O\varphi_t \rangle) > x \right] .$$

Kirkpatrick-R.-Schlein '20, R.-Seiringer '21: There exists, $C_{1,t}, C_{2,t} > 0$ s.t.

$$\limsup_{N \rightarrow \infty} \Lambda_{N,t}(x) \leq \frac{x^2}{2\sigma_t^2} + C_{1,t}x^3, \quad \liminf_{N \rightarrow \infty} \Lambda_{N,t}(x) \geq \frac{x^2}{2\sigma_t^2} - C_{2,t}x^{5/2}$$

► Variance σ_t^2 agrees with CLTs

► For $t > 0$ we have $C_{i,t} = e^{e^{Ct}}$



6. Large deviations


Characterized by

$$\Lambda_{N,t}(x) := -N^{-1} \log \mathbb{P}_{\psi_{N,t}} \left[N^{-1} \sum_{i=1}^N (Y_i^{N,t} - \langle \varphi_t, O \varphi_t \rangle) > x \right].$$

Kirkpatrick-R.-Schlein '20, R.-Seiringer '21: There exists, $C_{1,t}, C_{2,t} > 0$ s.t.

$$\limsup_{N \rightarrow \infty} \Lambda_{N,t}(x) \leq \frac{x^2}{2\sigma_t^2} + C_{1,t}x^3, \quad \liminf_{N \rightarrow \infty} \Lambda_{N,t}(x) \geq \frac{x^2}{2\sigma_t^2} - C_{2,t}x^{5/2}$$

- Variance σ_t^2 agrees with CLTs
- For $t > 0$ we have $C_{i,t} = e^{e^{Ct}}$
- For O such that $\|(-\Delta + 1)O(-\Delta + 1)^{-1}\|_{\text{op}} \leq C$



6. Large deviations


Characterized by

$$\Lambda_{N,t}(x) := -N^{-1} \log \mathbb{P}_{\psi_{N,t}} \left[N^{-1} \sum_{i=1}^N (Y_i^{N,t} - \langle \varphi_t, O \varphi_t \rangle) > x \right].$$

Kirkpatrick-R.-Schlein '20, R.-Seiringer '21: There exists, $C_{1,t}, C_{2,t} > 0$ s.t.

$$\limsup_{N \rightarrow \infty} \Lambda_{N,t}(x) \leq \frac{x^2}{2\sigma_t^2} + C_{1,t}x^3, \quad \liminf_{N \rightarrow \infty} \Lambda_{N,t}(x) \geq \frac{x^2}{2\sigma_t^2} - C_{2,t}x^{5/2}$$

- Variance σ_t^2 agrees with CLTs
- For $t > 0$ we have $C_{i,t} = e^{e^{Ct}}$
- For O such that $\|(-\Delta + 1)O(-\Delta + 1)^{-1}\|_{\text{op}} \leq C$
- For $\beta = 0$ only



Idea of the proof

Idea of the proof

Step 1: Generalization of Cramér's theorem,

Idea of the proof

Step 1: Generalization of Cramér's theorem,
LDE follow from:

Kirkpatrick-R.-Schlein ('21), R.-Seiringer ('22):

$$e^{N\left(\frac{\lambda^2}{2}\sigma_t^2 - C_t\lambda^3\right) - C\lambda} \leq \mathbb{E}_{\psi_{N,t}} \left[e^{\lambda \left(\sum_{i=1}^N Y_i^{N,t} - N\langle \varphi_t, O\varphi_t \rangle \right)} \right] \leq e^{N\left(\frac{\lambda^2}{2}\sigma_t^2 + C_t\lambda^3\right) + C_t\lambda}$$

Idea of the proof

Step 1: Generalization of Cramér's theorem,
LDE follow from:

Kirkpatrick-R.-Schlein ('21), R.-Seiringer ('22):

$$e^{N\left(\frac{\lambda^2}{2}\sigma_t^2 - C_t\lambda^3\right) - C_t\lambda} \leq \mathbb{E}_{\psi_{N,t}} \left[e^{\lambda \left(\sum_{i=1}^N Y_i^{N,t} - N\langle \varphi_t, O\varphi_t \rangle \right)} \right] \leq e^{N\left(\frac{\lambda^2}{2}\sigma_t^2 + C_t\lambda^3\right) + C_t\lambda}$$

Idea of the proof

Step 1: Generalization of Cramér's theorem,
LDE follow from:

Kirkpatrick-R.-Schlein ('21), R.-Seiringer ('22):

$$e^{N\left(\frac{\lambda^2}{2}\sigma_t^2 - C_t\lambda^3\right) - C_t\lambda} \leq \mathbb{E}_{\psi_{N,t}} \left[e^{\lambda \left(\sum_{i=1}^N Y_i^{N,t} - N\langle \varphi_t, O\varphi_t \rangle \right)} \right] \leq e^{N\left(\frac{\lambda^2}{2}\sigma_t^2 + C_t\lambda^3\right) + C_t\lambda}$$

Step 2: Proof of upper and lower bound.

Idea of Step 2

We study

$$\mathbb{E}_{\psi_{N,t}} \left[e^{\lambda \sum_{i=1}^N (Y_i^{N,t} - \langle \varphi_t, O \varphi_t \rangle)} \right] = \langle \psi_{N,t}, e^{\lambda Nd \Gamma(\check{O})} \psi_{N,t} \rangle$$

Idea of Step 2

We study

$$\begin{aligned}\mathbb{E}_{\psi_{N,t}} \left[e^{\lambda \sum_{i=1}^N (Y_i^{N,t} - \langle \varphi_t, O \varphi_t \rangle)} \right] &= \langle \psi_{N,t}, e^{\lambda Nd\Gamma(\tilde{O})} \psi_{N,t} \rangle \\ &= \langle \mathcal{W}_N(t; 0) \Omega, e^{\lambda \sqrt{N} \phi_+(q_t O \varphi_t) + \lambda d\Gamma(q_t \tilde{O} q_t)} \mathcal{W}_N(t; 0) \Omega \rangle\end{aligned}$$

Idea of Step 2

We study

$$\begin{aligned}\mathbb{E}_{\psi_{N,t}} \left[e^{\lambda \sum_{i=1}^N (Y_i^{N,t} - \langle \varphi_t, O \varphi_t \rangle)} \right] &= \langle \psi_{N,t}, e^{\lambda Nd\Gamma(\check{O})} \psi_{N,t} \rangle \\ &= \langle \mathcal{W}_N(t; 0) \Omega, e^{\lambda \sqrt{N} \phi_+ (q_t O \varphi_t) + \lambda d\Gamma(\cancel{q_t \check{O} q_t})} \mathcal{W}_N(t; 0) \Omega \rangle\end{aligned}$$

Idea of Step 2

We study

$$\begin{aligned}
 \mathbb{E}_{\psi_{N,t}} \left[e^{\lambda \sum_{i=1}^N (Y_i^{N,t} - \langle \varphi_t, O\varphi_t \rangle)} \right] &= \langle \psi_{N,t}, e^{\lambda Nd\Gamma(\tilde{O})} \psi_{N,t} \rangle \\
 &= \langle \mathcal{W}_N(t; 0)\Omega, e^{\lambda \sqrt{N}\phi_+(q_t O\varphi_t) + \lambda d\Gamma(q_t \tilde{O} q_t)} \mathcal{W}_N(t; 0)\Omega \rangle \\
 &\sim \langle \mathcal{W}_N(t; 0)\Omega, e^{\lambda \sqrt{N}\phi_+(q_t O\varphi_t)} \mathcal{W}_N(t; 0)\Omega \rangle
 \end{aligned}$$

Ingredients for the proof:

- Baker-Campbell Hausdorff formulas for non-commuting operators

Idea of Step 2

We study

$$\begin{aligned}
 \mathbb{E}_{\psi_{N,t}} \left[e^{\lambda \sum_{i=1}^N (Y_i^{N,t} - \langle \varphi_t, O \varphi_t \rangle)} \right] &= \langle \psi_{N,t}, e^{\lambda N d \Gamma(\tilde{O})} \psi_{N,t} \rangle \\
 &= \langle \mathcal{W}_N(t; 0) \Omega, e^{\lambda \sqrt{N} \phi_+ (q_t O \varphi_t) + \cancel{\lambda d \Gamma(q_t \tilde{O} q_t)}} \mathcal{W}_N(t; 0) \Omega \rangle \\
 &\sim \langle W_N(t; 0) \Omega, e^{\lambda \sqrt{N} \phi_+ (q_t O \varphi_t)} W_N(t; 0) \Omega \rangle \\
 &\sim \langle \mathcal{W}_\infty(t; 0) \Omega, e^{\lambda \sqrt{N} \phi_+ (q_t O \varphi_t)} \mathcal{W}_\infty(t; 0) \Omega \rangle
 \end{aligned}$$

Ingredients for the proof:

- Baker-Campbell Hausdorff formulas for non-commuting operators

Idea of Step 2

We study

$$\begin{aligned}
 \mathbb{E}_{\psi_{N,t}} \left[e^{\lambda \sum_{i=1}^N (Y_i^{N,t} - \langle \varphi_t, O \varphi_t \rangle)} \right] &= \langle \psi_{N,t}, e^{\lambda N d \Gamma(\tilde{O})} \psi_{N,t} \rangle \\
 &= \langle \mathcal{W}_N(t; 0) \Omega, e^{\lambda \sqrt{N} \phi_+ (q_t O \varphi_t) + \cancel{\lambda d \Gamma(q_t \tilde{O} q_t)}} \mathcal{W}_N(t; 0) \Omega \rangle \\
 &\sim \langle W_N(t; 0) \Omega, e^{\lambda \sqrt{N} \phi_+ (q_t O \varphi_t)} W_N(t; 0) \Omega \rangle \\
 &\sim \langle \mathcal{W}_\infty(t; 0) \Omega, e^{\lambda \sqrt{N} \phi_+ (q_t O \varphi_t)} \mathcal{W}_\infty(t; 0) \Omega \rangle \\
 &\sim \langle \Omega, e^{\lambda \sqrt{N} \phi_+ (\mathcal{L}_{(t;0)} q_t O \varphi_t)} \Omega \rangle
 \end{aligned}$$

Ingredients for the proof:

- ▶ Baker-Campbell Hausdorff formulas for non-commuting operators
- ▶ Exponential control of fluctuations by Gronwall type estimates

Idea of Step 2

We study

$$\begin{aligned}
 \mathbb{E}_{\psi_{N,t}} \left[e^{\lambda \sum_{i=1}^N (Y_i^{N,t} - \langle \varphi_t, O \varphi_t \rangle)} \right] &= \langle \psi_{N,t}, e^{\lambda N d \Gamma(\tilde{O})} \psi_{N,t} \rangle \\
 &= \langle \mathcal{W}_N(t; 0) \Omega, e^{\lambda \sqrt{N} \phi_+ (q_t O \varphi_t) + \cancel{\lambda d \Gamma(q_t \tilde{O} q_t)}} \mathcal{W}_N(t; 0) \Omega \rangle \\
 &\sim \langle \mathcal{W}_N(t; 0) \Omega, e^{\lambda \sqrt{N} \phi_+ (q_t O \varphi_t)} \mathcal{W}_N(t; 0) \Omega \rangle \\
 &\sim \langle \mathcal{W}_\infty(t; 0) \Omega, e^{\lambda \sqrt{N} \phi_+ (q_t O \varphi_t)} \mathcal{W}_\infty(t; 0) \Omega \rangle \\
 &\sim \langle \Omega, e^{\lambda \sqrt{N} \phi_+ (\mathcal{L}_{(t;0)} q_t O \varphi_t)} \Omega \rangle \\
 &\sim e^{N \lambda^2 \|\mathcal{L}_{(t;0)} q_t O \varphi_t\|_2^2}
 \end{aligned}$$

Ingredients for the proof:

- ▶ Baker-Campbell Hausdorff formulas for non-commuting operators
- ▶ Exponential control of fluctuations by Gronwall type estimates
- ▶ Control of $\mathcal{L}_{(t;0)}$

7. Quantum fluctuations in equilibrium

7. Quantum fluctuations in equilibrium

We consider N bosons on $\pi = [0, 1]^3$ described by

$$H_N = \sum_{i=1}^N (-\Delta_i) + \frac{1}{N} \sum_{1 \leq i < j \leq N} v_N^\beta(x_i - x_j)$$

with $v_N^\beta(x) = N^{3\beta} v(N^\beta x)$ and $\beta \in [0, 1]$.

7. Quantum fluctuations in equilibrium

We consider N bosons on $\pi = [0, 1]^3$ described by

$$H_N = \sum_{i=1}^N (-\Delta_i) + \frac{1}{N} \sum_{1 \leq i < j \leq N} v_N^\beta(x_i - x_j)$$

with $v_N^\beta(x) = N^{3\beta} v(N^\beta x)$ and $\beta \in [0, 1]$.

The ground state ψ_N exhibits Bose-Einstein condensation

$$\mathrm{Tr} \left| \gamma_{\psi_N}^{(1)} - N |\varphi\rangle \langle \varphi| \right| \leq C, \quad \text{for all } N \in \mathbb{N},$$

following from **Lieb-Seiringer (2001)** for all $\beta \in [0, 1]$

7. Quantum fluctuations in equilibrium

We consider N bosons on $\pi = [0, 1]^3$ described by

$$H_N = \sum_{i=1}^N (-\Delta_i) + \frac{1}{N} \sum_{1 \leq i < j \leq N} v_N^\beta(x_i - x_j)$$

with $v_N^\beta(x) = N^{3\beta} v(N^\beta x)$ and $\beta \in [0, 1]$.

The ground state ψ_N exhibits Bose-Einstein condensation

$$\mathrm{Tr} \left| \gamma_{\psi_N}^{(1)} - N |\varphi\rangle\langle\varphi| \right| \leq C, \quad \text{for all } N \in \mathbb{N}, \quad \text{But: } \psi_N \approx \varphi^{\otimes N}$$

following from **Lieb-Seiringer (2001)** for all $\beta \in [0, 1]$

See: **Brennecke, Cenatiempo, Lewin, Nam, Schlein, Serfaty, Solovej,...**

7. Quantum fluctuations in equilibrium

We consider N bosons on $\pi = [0, 1]^3$ described by

$$H_N = \sum_{i=1}^N (-\Delta_i) + \frac{1}{N} \sum_{1 \leq i < j \leq N} v_N^\beta(x_i - x_j)$$

with $v_N^\beta(x) = N^{3\beta} v(N^\beta x)$ and $\beta \in [0, 1]$.

The ground state ψ_N exhibits Bose-Einstein condensation

$$\mathrm{Tr} \left| \gamma_{\psi_N}^{(1)} - N |\varphi\rangle\langle\varphi| \right| \leq C, \quad \text{for all } N \in \mathbb{N}, \quad \text{But: } \psi_N \approx \varphi^{\otimes N}$$

following from **Lieb-Seiringer (2001)** for all $\beta \in [0, 1]$

See: **Brennecke, Cenatiempo, Lewin, Nam, Schlein, Serfaty, Solovej,...**

Random variables Y_i^N : with law

$$\mathbb{P}_{\psi_N} [Y_i^N \in A] = \langle \psi_N, \mathbb{1}_A(O^{(i)}) \psi_N \rangle$$

for bounded self-adjoint operator O are **identically distributed** and **correlated**.

7.1 Limit theorems

7.1 Limit theorems

Law of large numbers: follows from BEC

$$\lim_{N \rightarrow \infty} \mathbb{P}_{\psi_N} \left[\left| \frac{1}{N} \sum_{i=1}^N (Y_i^N - \langle \varphi, O\varphi \rangle) \right| > \delta \right] = 0$$

for any $\delta > 0$ and appropriate $\varphi \in L^2(\pi)$ for all $\beta \in [0, 1]$.

7.1 Limit theorems

Law of large numbers: follows from BEC

$$\lim_{N \rightarrow \infty} \mathbb{P}_{\psi_N} \left[\left| \frac{1}{N} \sum_{i=1}^N (Y_i^N - \langle \varphi, O\varphi \rangle) \right| > \delta \right] = 0$$

for any $\delta > 0$ and appropriate $\varphi \in L^2(\pi)$ for all $\beta \in [0, 1]$.

7.1 Limit theorems

Law of large numbers: follows from BEC

$$\lim_{N \rightarrow \infty} \mathbb{P}_{\psi_N} \left[\left| \frac{1}{N} \sum_{i=1}^N (Y_i^N - \langle \varphi, O\varphi \rangle) \right| > \delta \right] = 0$$

for any $\delta > 0$ and appropriate $\varphi \in L^2(\pi)$ for all $\beta \in [0, 1]$.

Central limit theorem proven by **R.-Schlein (2019)**

$$\lim_{N \rightarrow \infty} \mathbb{P}_{\psi_N} \left[\frac{1}{\sqrt{N}} \sum_{i=1}^N (Y_i^N - \langle \varphi, O\varphi \rangle) \in [a, b] \right] = \mathbb{P} \left[\mathcal{G} \in [a, b] \right]$$

7.1 Limit theorems

Law of large numbers: follows from BEC

$$\lim_{N \rightarrow \infty} \mathbb{P}_{\psi_N} \left[\left| \frac{1}{N} \sum_{i=1}^N (Y_i^N - \langle \varphi, O\varphi \rangle) \right| > \delta \right] = 0$$

for any $\delta > 0$ and appropriate $\varphi \in L^2(\pi)$ for all $\beta \in [0, 1]$.

Central limit theorem proven by **R.-Schlein (2019)**

$$\lim_{N \rightarrow \infty} \mathbb{P}_{\psi_N} \left[\frac{1}{\sqrt{N}} \sum_{i=1}^N (Y_i^N - \langle \varphi, O\varphi \rangle) \in [a, b] \right] = \mathbb{P} \left[\mathcal{G} \in [a, b] \right]$$

and \mathcal{G} is the **centered Gaussian random variable** with variance $\|\sigma\|_2^2$

7.1 Limit theorems

Law of large numbers: follows from BEC

$$\lim_{N \rightarrow \infty} \mathbb{P}_{\psi_N} \left[\left| \frac{1}{N} \sum_{i=1}^N (Y_i^N - \langle \varphi, O\varphi \rangle) \right| > \delta \right] = 0$$

for any $\delta > 0$ and appropriate $\varphi \in L^2(\pi)$ for all $\beta \in [0, 1]$.

Central limit theorem proven by **R.-Schlein (2019)**

$$\lim_{N \rightarrow \infty} \mathbb{P}_{\psi_N} \left[\frac{1}{\sqrt{N}} \sum_{i=1}^N (Y_i^N - \langle \varphi, O\varphi \rangle) \in [a, b] \right] = \mathbb{P} \left[\mathcal{G} \in [a, b] \right]$$

and \mathcal{G} is the **centered Gaussian random variable** with variance $\|\sigma\|_2^2$ given by

$$\sigma_p = \cosh(\mu_p) \widehat{(qO\varphi)}(p) \sinh(\mu_p) \widehat{(qO\varphi)}(p), \quad \text{for all } p \in (2\pi\mathbb{Z})^3$$

and with

$$\mu_p = \frac{1}{4} \log \left(\frac{p^2}{p^2 + 2\widehat{v}(p)} \right), \quad \text{for } \beta = 0$$

7.1 Limit theorems

Law of large numbers: follows from BEC

$$\lim_{N \rightarrow \infty} \mathbb{P}_{\psi_N} \left[\left| \frac{1}{N} \sum_{i=1}^N (Y_i^N - \langle \varphi, O\varphi \rangle) \right| > \delta \right] = 0$$

for any $\delta > 0$ and appropriate $\varphi \in L^2(\pi)$ for all $\beta \in [0, 1]$.

Central limit theorem proven by **R.-Schlein (2019)**

$$\lim_{N \rightarrow \infty} \mathbb{P}_{\psi_N} \left[\frac{1}{\sqrt{N}} \sum_{i=1}^N (Y_i^N - \langle \varphi, O\varphi \rangle) \in [a, b] \right] = \mathbb{P} \left[\mathcal{G} \in [a, b] \right]$$

and \mathcal{G} is the **centered Gaussian random variable** with variance $\|\sigma\|_2^2$ given by

$$\sigma_p = \cosh(\mu_p) \widehat{(qO\varphi)}(p) \sinh(\mu_p) \widehat{(qO\varphi)}(p), \quad \text{for all } p \in (2\pi\mathbb{Z})^3$$

and with

$$\mu_p = \frac{1}{4} \log \left(\frac{p^2}{p^2 + 16\pi a_0} \right), \quad \text{for } \beta = 1$$

7.2 Large deviations

Characterized by

$$\Lambda_{\psi_N}(x) := -N^{-1} \log \mathbb{P}_{\psi_N} \left[N^{-1} \sum_{i=1}^N (Y_i^N - \langle \varphi, O\varphi \rangle) > x \right] .$$

7.2 Large deviations

Characterized by

$$\Lambda_{\psi_N}(x) := -N^{-1} \log \mathbb{P}_{\psi_N} \left[N^{-1} \sum_{i=1}^N (Y_i^N - \langle \varphi, O\varphi \rangle) > x \right] .$$

R '23: There exists, $C_1, C_2 > 0$ such that

$$\limsup_{N \rightarrow \infty} \Lambda_{\psi_N}(x) \leq \frac{x^2}{2\sigma^2} + C_1 x^3, \quad \liminf_{N \rightarrow \infty} \Lambda_{\psi_N}(x) \geq \frac{x^2}{2\sigma^2} - C_2 x^{5/2}$$

7.2 Large deviations

Characterized by

$$\Lambda_{\psi_N}(x) := -N^{-1} \log \mathbb{P}_{\psi_N} \left[N^{-1} \sum_{i=1}^N (Y_i^N - \langle \varphi, O\varphi \rangle) > x \right] .$$

R '23: There exists, $C_1, C_2 > 0$ such that

$$\limsup_{N \rightarrow \infty} \Lambda_{\psi_N}(x) \leq \frac{x^2}{2\sigma^2} + C_1 x^3, \quad \liminf_{N \rightarrow \infty} \Lambda_{\psi_N}(x) \geq \frac{x^2}{2\sigma^2} - C_2 x^{5/2}$$

► Variance σ^2 agrees with CLTs

7.2 Large deviations

Characterized by

$$\Lambda_{\psi_N}(x) := -N^{-1} \log \mathbb{P}_{\psi_N} \left[N^{-1} \sum_{i=1}^N (Y_i^N - \langle \varphi, O\varphi \rangle) > x \right] .$$

R '23: There exists, $C_1, C_2 > 0$ such that

$$\limsup_{N \rightarrow \infty} \Lambda_{\psi_N}(x) \leq \frac{x^2}{2\sigma^2} + C_1 x^3, \quad \liminf_{N \rightarrow \infty} \Lambda_{\psi_N}(x) \geq \frac{x^2}{2\sigma^2} - C_2 x^{5/2}$$

- Variance σ^2 agrees with CLTs
- For O such that $\|(-\Delta + 1)O(-\Delta + 1)^{-1}\|_{\text{op}} \leq C$

7.2 Large deviations

Characterized by

$$\Lambda_{\psi_N}(x) := -N^{-1} \log \mathbb{P}_{\psi_N} \left[N^{-1} \sum_{i=1}^N (Y_i^N - \langle \varphi, O\varphi \rangle) > x \right] .$$

R '23: There exists, $C_1, C_2 > 0$ such that

$$\limsup_{N \rightarrow \infty} \Lambda_{\psi_N}(x) \leq \frac{x^2}{2\sigma^2} + C_1 x^3, \quad \liminf_{N \rightarrow \infty} \Lambda_{\psi_N}(x) \geq \frac{x^2}{2\sigma^2} - C_2 x^{5/2}$$

- Variance σ^2 agrees with CLTs
- For O such that $\|(-\Delta + 1)O(-\Delta + 1)^{-1}\|_{\text{op}} \leq C$
- For $\beta = 0$ only

7.2 Large deviations

Characterized by

$$\Lambda_{\psi_N}(x) := -N^{-1} \log \mathbb{P}_{\psi_N} \left[N^{-1} \sum_{i=1}^N (Y_i^N - \langle \varphi, O\varphi \rangle) > x \right] .$$

R '23: There exists, $C_1, C_2 > 0$ such that

$$\limsup_{N \rightarrow \infty} \Lambda_{\psi_N}(x) \leq \frac{x^2}{2\sigma^2} + C_1 x^3, \quad \liminf_{N \rightarrow \infty} \Lambda_{\psi_N}(x) \geq \frac{x^2}{2\sigma^2} - C_2 x^{5/2}$$

- Variance σ^2 agrees with CLTs
- For O such that $\|(-\Delta + 1)O(-\Delta + 1)^{-1}\|_{\text{op}} \leq C$
- For $\beta = 0$ only

For $O = qOq$: and $\beta = 1$, it holds **R-Nam '23**

$$-\limsup_{N \rightarrow \infty} \log \mathbb{P}_{\psi_N} \left[\sum_{i=1}^N Y_i^N - \mu_0 > x \right] \leq \frac{x^2}{\sigma_0^2} + Cx^3 .$$

7.3 Empirical measures

7.3 Empirical measures

For any $A \subset \mathcal{B}(\mathbb{R})$ compare

$$\nu_N(A) := \frac{1}{N} \sum_{i=1}^N \delta_{Y_i^N}(A) \quad \text{and} \quad \nu_\varphi(A) := \langle \varphi, \mathbb{1}_A(O) \varphi \rangle_{L^2(\pi)} .$$

using **Wasserstein metric** \mathcal{W}_1 .

7.3 Empirical measures

For any $A \subset \mathcal{B}(\mathbb{R})$ compare

$$\nu_N(A) := \frac{1}{N} \sum_{i=1}^N \delta_{Y_i^N}(A) \quad \text{and} \quad \nu_\varphi(A) := \langle \varphi, \mathbb{1}_A(O) \varphi \rangle_{L^2(\pi)}.$$

using **Wasserstein metric** \mathcal{W}_1 .

Portinale-R.-Virostek '23: Law of large numbers

$$\lim_{N \rightarrow \infty} \mathbb{P} \left[|\mathcal{W}_1(\nu_N, \nu_\varphi)| > \delta \right] = 0.$$

7.3 Empirical measures

For any $A \subset \mathcal{B}(\mathbb{R})$ compare

$$\nu_N(A) := \frac{1}{N} \sum_{i=1}^N \delta_{Y_i^N}(A) \quad \text{and} \quad \nu_\varphi(A) := \langle \varphi, \mathbb{1}_A(O) \varphi \rangle_{L^2(\pi)}.$$

using **Wasserstein metric** \mathcal{W}_1 .

Portinale-R.-Virostek '23: Law of large numbers

$$\lim_{N \rightarrow \infty} \mathbb{P} \left[|\mathcal{W}_1(\nu_N, \nu_\varphi)| > \delta \right] = 0.$$

Note:

► Proof for $\beta = 1$

7.3 Empirical measures

For any $A \subset \mathcal{B}(\mathbb{R})$ compare

$$\nu_N(A) := \frac{1}{N} \sum_{i=1}^N \delta_{Y_i^N}(A) \quad \text{and} \quad \nu_\varphi(A) := \langle \varphi, \mathbb{1}_A(O) \varphi \rangle_{L^2(\pi)}.$$

using **Wasserstein metric** \mathcal{W}_1 .

Portinale-R.-Virostek '23: Law of large numbers

$$\lim_{N \rightarrow \infty} \mathbb{P}[|\mathcal{W}_1(\nu_N, \nu_\varphi)| > \delta] = 0.$$

Note:

- ▶ Proof for $\beta = 1$
- ▶ Gaussian fluctuations around ν_φ

8. Open questions

8. Open questions

8. Open questions

- ▶ Sanov's type estimates ?

8. Open questions

- ▶ Sanov's type estimates ?
- ▶ Large deviation estimates for $\beta \in (0, 1]$

8. Open questions

- ▶ Sanov's type estimates ?
- ▶ Large deviation estimates for $\beta \in (0, 1]$
- ▶ Higher order terms of rate function for all $\beta \in [0, 1]$

8. Open questions

- ▶ Sanov's type estimates ?
- ▶ Large deviation estimates for $\beta \in (0, 1]$
- ▶ Higher order terms of rate function for all $\beta \in [0, 1]$
- ▶ Large deviation principle?

8. Open questions

- ▶ Sanov's type estimates ?
- ▶ Large deviation estimates for $\beta \in (0, 1]$
- ▶ Higher order terms of rate function for all $\beta \in [0, 1]$
- ▶ Large deviation principle?
- ▶ Positive Temperatures?

8. Open questions

- ▶ Sanov's type estimates ?
- ▶ Large deviation estimates for $\beta \in (0, 1]$
- ▶ Higher order terms of rate function for all $\beta \in [0, 1]$
- ▶ Large deviation principle?
- ▶ Positive Temperatures?
- ▶ Thermodynamic limit?

8. Open questions

- ▶ Sanov's type estimates ?
- ▶ Large deviation estimates for $\beta \in (0, 1]$
- ▶ Higher order terms of rate function for all $\beta \in [0, 1]$
- ▶ Large deviation principle?
- ▶ Positive Temperatures?
- ▶ Thermodynamic limit?

Thanks for your attention.